

Four-dimensional effective theory of gravity embeddable in a distorted Randall-Sundrum brane scenario

I. Quiros*, R. Bonal[†] and R. Cardenas[‡]

Dpto. Fisica. Universidad Central de Las Villas. Santa Clara CP 54830. Villa Clara. Cuba

(February 7, 2008)

Abstract

We present an effective four-dimensional formulation of the laws of gravity that respects the main features of a higher (five)-dimensional scenario of Randall-Sundrum type. The geometrical structure of the theory is that of a Weyl-integrable configuration. Standard general relativity over Riemann geometry is recovered through breaking of conformal symmetry. The singularity problem is treated. The local problem in the case of the static, spherically-symmetric Schwarzschild metric and the cosmological issue, in the case of Friedmann-Robertson-Walker perfect fluid filled universe, are treated separately. Vanishing of the spacetime singularities for some values of the free parameter of the theory is achieved. The implications of the results obtained for brane stabilization in the higher-dimensional structure are briefly discussed.

Typeset using REVTeX

*israel@uclv.etecsa.cu

[†]bonal@uclv.etecsa.cu

[‡]rcardenas@uclv.etecsa.cu

I. INTRODUCTION

There are two main ingredients that enter the basis of the unification scheme of the fundamental interactions. First, multiple dimensions are required. In fact, modern field theory suggests spacetime to be $4 + n$ -dimensional ($n = 6$ for string theory and $n = 7$ for supergravity). It is usually assumed that the additional dimensions may be compactified down to size of the order the Planck length or below so they are not observable. However, recent developments suggest that some of the additional dimensions may have compactification radii larger than the Planck length without conflict with observations [1]. Besides, it is possible that gravity remains confined to a four-dimensional slice of the bulk spacetime leading to extra-dimensions being even infinite in extent [2]. The above scenarios with large extra-dimensions start from the assumption that ordinary matter fields (in particular standard model particles) are confined to a three-brane (four spacetime dimensions) embedded in a higher-dimensional bulk spacetime. These models provide, in particular, successful resolutions of the mass hierarchy problem. Even if none of these scenarios with large extra-dimensions correctly describe our present universe, their potential for describing the early universe is obvious since there is strong evidence that the early universe underwent a phase where it was five-dimensional [3–5].

The second basic ingredient of the unification scheme is based on the assumption that the universe must evolve from maximally symmetric states into states with broken symmetries. The aim of the symmetry breaking mechanism is always to realize a transition from totally symmetric spacetimes (for instance de Sitter), where the matter content is due to quantum fields in a polarized vacuum state, to our asymmetric universe (for instance Friedmann-Robertson-Walker) with four fundamental interactions and the known families of particles. One of the basic symmetries that may be linked with gravitation and that must be broken at present is conformal (Weyl) symmetry. It consists of a conformal rescaling of the spacetime metric and a field redefinition. There are some interesting attempts for explaining, for instance, the origin of the gravitational coupling of matter [6] and the mass generation [7]

by breaking of the original conformal symmetry of the gravitational theory.

Following the above arguments coming from the unification scheme one may argue that, although Einstein's general relativity is the leading theory of gravity correctly explaining current experimental observations at present epoch in the evolution of the universe, in the distant past, when some currently broken symmetries were exact symmetries of the laws of physics (including gravity), some other effective four-dimensional theory of gravity with higher symmetry was the leading one (for instance low-energy string theory). The more important constraint on this higher-symmetric theory is that general relativity must be obtained from it by breaking of the additional symmetries. The same line of reasoning may be applied to geometry. General relativity is built over Riemann geometry. However, we may argue that, in the distant past, some other higher-symmetric geometry was the leading one. Riemann geometry is, in this context, a symmetry-broken state of the higher-symmetric geometry.

Thinking along these lines one may conjecture that the minimal requirement any "geometrically consistent" four-dimensional description of the laws of gravity should fulfill is that the effective action for the theory and the geometry over which it is built must share the same symmetries. For the sake of brevity we shall call this as "geometric consistency conjecture". This conjecture may appear naive, however it rules out, for instance, Brans-Dicke (BD) theory in its usual formulation [8]. In fact, the pure gravitational part for BD theory $S = \int d^4x \sqrt{-g} e^{-\psi} (R - \omega (\nabla \psi)^2)$, where R is the Ricci scalar, ψ is the BD field and ω is the BD coupling parameter, is conformally-invariant. Even the presence of radiation matter fields does not break the conformal invariance of the theory (see for instance Ref. [5] and references therein). Nevertheless, BD theory is built over Riemannian spacetimes that do not possess the conformal symmetry.

Weyl-integrable geometry (WIG), instead, respects conformal (Weyl) invariance. It is, precisely, its distinctive feature. Moreover, variation *a la* Palatini of the above effective action for Brans-Dicke theory, leads *a posteriori* to spacetimes of Weyl-integrable configuration as the basic geometric structure of the theory rather than Riemannian spacetimes (see, for

instance, Ref. [9]). Therefore a kind of Brans-Dicke theory over Weyl-integrable geometry leads to "geometrical consistency" being recovered and, at the same time, is more in the spirit of the unification scheme of the fundamental interactions.

Usually, higher-dimensional and higher-symmetric theories are built by postulating the higher-dimensional equations for the theory. From this higher-dimensional formulation four-dimensional physics is recovered by using different procedures. One example is five-dimensional Kaluza-Klein theory where the compactification procedure is used. Recovering of four-dimensional physics in other alternatives as, for instance, the recently proposed scenario of Randall and Sundrum with a non-compact extra-dimension [2], relies on the confining of the matter degrees of freedom on a four-dimensional slice (brane) of the bulk spacetime. These higher-dimensional descriptions are, in general, difficult to manage since, in most cases, complicated junction conditions should be imposed on the fields that are present in the theory [10].

The present paper is aimed at a derivation of a four-dimensional effective theory of gravity that should be embeddable in a five-dimensional "distorted" Randall-Sundrum brane scenario. Unlike the standard approach starting by postulating from the beginning the higher-dimensional equations of the theory, in this paper we shall develop an alternative approach that is based on an essentially four-dimensional way of thinking. We start with the Randall-Sundrum scenario improved in Ref. [11] and then we "distort" it by relaxing some requirements as, for instance, orbifold symmetry and Poincare invariance. Then we elaborate on this "distorted" five-dimensional brane scenario by studying the effective geometry induced on the four-dimensional manifold. It results in a Weyl-integrable geometry that may be seen as a distorted Riemann geometry. The next step is to postulate a four-dimensional theory of gravity that may be embedded in the higher-symmetric structure, i.e., a theory sharing some of the basic properties of the "distorted" brane set-up worked out previously. For instance, the theory we search for should be built over Weyl-integrable geometry, then it should be Weyl-symmetric. Besides, the matter degrees of freedom of the theory should be coupled to the metric induced on the "visible" brane of the higher-dimensional scenario in-

stead of the four-dimensional "Planck" metric. There should be in the theory some "signal" from the extra-dimension, etc.

The resulting four-dimensional effective formulation of the laws of gravity is intended to describe physics over a time scale extending from (approximately) the Planck era to the conformal symmetry broken era. Once conformal (Weyl) symmetry is broken, four-dimensional general relativity and, consequently, Riemann geometry over which it is built, are recovered from the present description. In other words, general relativity over Riemann geometry must be seen as a Weyl symmetry broken state of the effective formulation presented in this paper.

Although Weyl symmetry at present is broken on cosmological scales, it may be argued that, near of gigantic concentrations of energy where the four-dimensional slice (brane) which we live in may be considerably distorted or bent in the extra-dimension, Weyl symmetry should be an exact symmetry of the laws of physics. Consequently our brane-inspired four-dimensional formulation may be relevant for the description of the physics near of such gigantic concentrations of energy. In this sense the black hole issue is of particular interest and will be treated here.

Finally it should be remarked that, since the present four-dimensional effective theory is embeddable in a higher-dimensional scenario, solving the equations of the theory would yield better understanding of the higher-dimensional structure.

The paper has been organized as follows. In Sec. II we consider a five-dimensional brane scenario that can be viewed as a distorted Randall-Sundrum scenario. In Sec. III we elaborate on this distorted brane set-up and we shall show that the four-dimensional effective geometrical structure it yields is one of Weyl-integrable configuration. In Sec. IV we derive a four-dimensional effective theory that is embeddable in the higher-dimensional brane scenario of Sec. II. The singularity problem is treated in Sec. V within the frame of this effective description of gravity. The local problem in the case of the static, spherically-symmetric Schwarzschild metric and the cosmological issue, in the case of Friedmann-Robertson-Walker perfect fluid filled universe, are treated separately. Vanishing of the spacetime singularities

for some values of the free parameter of the theory is very encouraging. The implications of the results obtained for the higher-dimensional structure are briefly discussed, in particular for the problem of brane stabilization. Conclusions are given in Sec. VI.

II. DISTORTED RANDALL-SUNDRUM SCENARIO

As a model for a higher-dimensional and higher-symmetric structure in which the desired effective four-dimensional theory of gravity is to be "embedded", we shall choose a five-dimensional brane spacetime. Being more specific; we shall study a Randall-Sundrum scenario [2] improved in Ref. [11].

In the original Randall-Sundrum-Lykken (RSL) scenario of Ref. [11] the graviton zero mode is confined on one of the branes (the "Planck" brane). The other brane is located a given distance away from this one in the additional y -direction. The "visible" matter degrees of freedom (the standard model matter fields) are supposed to be trapped there. Consequently this second brane is acknowledged as the "visible" brane.

The basic ingredient of the RSL scenario is a non-factorizable metric

$$ds^2 = e^{-2ky} \eta_{mn} dx^n dx^m + dy^2, \quad (2.1)$$

where η_{ab} is the flat Minkowski metric, $x \equiv x^a$ accounts for the usual four-dimensional coordinates and the extra coordinate $y \in [0, \pi r_c]$, r_c being the "compactification radius". Put in simple words: the four-dimensional metric is multiplied by a warp factor. In this model orbifold symmetry is imposed upon the y -direction ($y \rightarrow -y$), i.e., it is required to work on the space $S^1/\mathbf{Z}_2$. In addition four-dimensional Poincare invariance is also satisfied. The RSL scenario relies on the assumption that the branes are already stabilized in the sense that their separation in the y -direction is not dependent on the time-coordinate. However, in general, the branes are in relative motion so it is necessary to choose a reference brane [12]. Take, for instance, the Planck brane to be the reference brane. Therefore, in respect to a chosen normal Gaussian coordinate system the y -coordinate of the "visible" brane is time-dependent $y = y(t)$ [12]. The following reasoning allows generalizing of the former procedure

to include dependence on the spacetime coordinates x in general: $y = y(x)$. A key-point in standard four-dimensional theories of gravity is that any matter distribution causes the spacetime around it to become curved. This statement is true independent of the dimension of the spacetime. Consequently, any matter distribution, no matter whether it lies on a four-dimensional slice (hypersurface) or in the bulk spacetime, causes the five-dimensional (bulk) spacetime to become bent around it [13]. This yields that the y -direction is bent by the effect of the standard model matter fields living on the "visible" brane. Therefore, we may specify the Planck (reference) brane to be located at $y = 0$ and, in respect to it, we may pick a normal Gaussian coordinate system such that the following anzats for the five-dimensional line-element takes place:

$$ds^2 = e^{-2ky(x)} g_{nm} dx^n dx^m + dy^2(x). \quad (2.2)$$

In respect to this reference system the curvature effect produced by the standard model matter trapped on the "visible" brane is viewed as local distortions of this brane from flat configuration. In fact, the separation of the "visible" brane from the Planck one depends now on the four-dimensional spacetime coordinates.

The five-dimensional metric of the bulk spacetime has the components $G_{ab} = e^{-2ky(x)} g_{ab}(x)$ and $G_{44} = 1$. We see that the metric induced on the Planck brane $G_{ab}(x, y = 0)$ coincides with the four-dimensional metric g_{ab} , while the metric induced on the "visible" brane is $\bar{g}_{ab} = G_{ab}(x, y = \frac{1}{2k}\psi_0(x)) = e^{-\psi_0(x)} g_{ab}$. For the matter that is trapped in this brane it is required that $dy = \frac{1}{2k}d\psi_0(x) = 0$, i.e., $\psi_{0,n}dx^n = 0$. However, in what follows we shall relax this condition by supposing that ψ_0 may be, also, a function of additional internal degrees of freedom σ_i coming, for instance, from additional compactified extra-dimensions. Therefore $d\psi_0 = 0$ on the "visible" brane yields $-(\partial\psi_0/\partial\sigma_i)d\sigma_i = \psi_{0,n}dx^n$ so that, in principle, $\psi_{0,n}dx^n \neq 0$. The need for this requirement will be evident in what follows.

In the present scenario additional "distortion" of the original Randall-Sundrum set-up is achieved by dropping of the orbifold symmetry and Poincare invariance. The resulting non-factorizable five-dimensional line-element with curved extra-direction ψ may written as

$$ds^2 = e^{-2\psi(x)} g_{nm} dx^n dx^m + \frac{1}{4k^2} d\psi^2(x). \quad (2.3)$$

Requiring invariance of this line-element in respect to the following coordinate transformation

$$\begin{aligned} x^a &\rightarrow x^a, \\ \psi &\rightarrow \psi + 2 \ln \Omega, \end{aligned} \quad (2.4)$$

where $\Omega(x)$ is a non-vanishing smooth function, yields that the four-dimensional metric g_{ab} and the constant factor $\frac{1}{4k^2}$ should be transformed according to

$$g_{ab} \rightarrow \Omega^2 g_{ab}, \quad (2.5)$$

and

$$1/4k^2 \rightarrow \frac{1/4k^2}{(1 + 2\Omega^{-1}\Omega_{,\psi})^2}, \quad (2.6)$$

respectively. In other words, the coordinate transformation (2.4) in the bulk spacetime, induces a conformal transformation of the four-dimensional metric together with a transformation of the free parameter (Eq. (2.6)). This fact will be exploited in what follows.

III. EFFECTIVE FOUR-DIMENSIONAL WEYL-INTEGRABLE GEOMETRY

Irrespective of the concrete theory of gravity one chooses it is usually assumed that the five-dimensional spacetime has Riemann configuration. It is, precisely, the starting point of the reasoning-line to be developed here.

Riemann structure of the bulk spacetime means vanishing of the covariant derivative of the five-dimensional metric

$$G_{AB;C} = 0. \quad (3.1)$$

Condition (3.1) yields, in turn, that the affine connection of the bulk coincides with the Christoffel symbols of the five-dimensional metric:

$$\{ \begin{smallmatrix} A \\ BC \end{smallmatrix} \} = \frac{1}{2} G^{AN} (G_{BN,C} + G_{CN,B} - G_{BC,N}). \quad (3.2)$$

Matter particles would follow geodesic curves of this geometry:

$$\frac{d^2 x^A}{ds^2} + \{ \begin{smallmatrix} A \\ MN \end{smallmatrix} \} \frac{dx^M}{ds} \frac{dx^N}{ds} = 0. \quad (3.3)$$

In the "distorted" Randall-Sundrum scenario proposed here (line-element (2.3)), the standard model matter particles are confined on the "visible" brane obeying the surface equation $y = \psi(x)$, such that $d\psi = 0$. Consequently, the set of equations (3.3) reduces to

$$\frac{d^2 x^a}{d\bar{s}^2} + \{ \begin{smallmatrix} a \\ mn \end{smallmatrix} \}_4 \frac{dx^m}{d\bar{s}} \frac{dx^n}{d\bar{s}} = 0, \quad (3.4)$$

where $\{ \begin{smallmatrix} a \\ bc \end{smallmatrix} \}_4$ is the four-dimensional part of the Christoffel symbols (3.2) and $d\bar{s}^2 = e^{-\psi} g_{mn} dx^m dx^n = e^{-\psi} ds_4^2$. In terms of the four-dimensional metric g_{ab} , Eq. (3.4) may be written as

$$\frac{d^2 x^a}{ds_4^2} + \frac{1}{2} \psi_{,n} \frac{dx^n}{ds_4} \frac{dx^a}{ds_4} + \{ \begin{smallmatrix} a \\ mn \end{smallmatrix} \} \frac{dx^m}{ds_4} \frac{dx^n}{ds_4} - \psi_{,n} \left(\frac{dx^n}{ds_4} \frac{dx^a}{ds_4} - \frac{1}{2} g^{na} \right) = 0, \quad (3.5)$$

where $\{ \begin{smallmatrix} a \\ bc \end{smallmatrix} \}$ are the Christoffel symbols of the four-dimensional metric g_{ab} .

At first sight, it seems that standard matter particles do not follow geodesics of the four-dimensional geometry. However this is not true. In fact, note that condition (3.1) for the five-dimensional metric can be reduced, in the present scenario, to the following condition for the four-dimensional one¹

$$g_{ab;c} = \psi_{,c} g_{ab}, \quad (3.6)$$

where semicolon denotes covariant differentiation in a general affine sense. Consequently, the affine connection of the four-dimensional "effective" manifold coincide with the Weyl affine connection

$$\Gamma_{bc}^a = \{ \begin{smallmatrix} a \\ bc \end{smallmatrix} \} - \frac{1}{2} (\psi_{,b} \delta_c^a + \psi_{,c} \delta_b^a - g_{bc} \psi^{,a}), \quad (3.7)$$

¹The other equalities contained in Eq. (3.1) are now just identities

where the Weyl gauge vector is the gradient of the curved extra-coordinate $\psi_{,a}$. This yields that the "effective" four-dimensional geometry is of Weyl-integrable configuration. Consequently, the units of measure of the "effective" geometry changes length along transport, i.e., they depend on location on the four-dimensional manifold. Hence, one should expect that mass units should be, indeed, functions of the four-dimensional coordinates too. In particular, if the standard model particle's mass is given through $m = e^{\frac{\psi}{2}} m_0$, where m_0 is a constant and, by defining the four-momentum of the particle through $P^a = m \frac{dx^a}{ds_4}$, then Eq. (3.5) may be put in the form

$$\frac{dP^a}{ds_4} + m \Gamma_{mn}^a \frac{dx^m}{ds_4} \frac{dx^n}{ds_4} = 0, \quad (3.8)$$

where, as before, Γ_{bc}^a is the affine connection of the Weyl-integrable four-dimensional manifold with metric tensor g_{ab} (see Eq. (3.7)). Therefore, standard model matter particles confined on the "visible" brane follow geodesics in a four-dimensional manifold of Weyl-integrable "effective" configuration and their "effective" masses are functions of the four-dimensional coordinates through the curved extra-dimension $\psi(x)$.² It is straightforward noting that the resulting Weyl-integrable "effective" geometric structure is Weyl-symmetric, i.e., it is invariant under the Weyl (gauge) transformations (2.4) and (2.5). Now the requirement that the curved extra-direction ψ should depend on additional internal degrees of freedom becomes clear. In fact, suppose that the contrary condition holds, i.e., there are not internal degrees of freedom so that $\psi_{,n} dx^n = 0$. In this case Eq. (3.8) simplifies to

$$\frac{dP^a}{ds_4} + m \{^a_{mn}\} \frac{dx^m}{ds_4} \frac{dx^n}{ds_4} = 0, \quad (3.9)$$

that is not Weyl-symmetric in the sense described above. Therefore, if one is looking for a Weyl-symmetric "geometrically consistent" effective theory, the requirement $\psi_{,n} dx^n \neq 0$ should be fulfilled. This together with the condition that, on a given brane $d\psi = 0$, leads to ψ being function of additional internal degrees of freedom.

²The fact that variable masses may be linked with extra-dimensions is studied, for instance, in Ref. [14]

As in the original RSL scenario, in the present set-up, the brane where the standard model matter fields are trapped can be regarded as a probe of the geometry determined by the Planck brane [11].

IV. WEYL-SYMMETRIC EFFECTIVE FOUR-DIMENSIONAL THEORY OF GRAVITY

We now elaborate on the distorted RSL scenario presented in Sec. II in order to search for an effective four-dimensional theory of gravity consistent with higher (five)-dimensional and higher (Weyl)-symmetric spacetime structure.

The theory we are searching for should reflect invariance under transformations (2.4-2.6). Besides, following the conjecture of "geometrical consistency" presented in Sec. I, it should be built over an effective Weyl-integrable geometry given by the law (3.6). Moreover, the theory should be "embeddable" in the "distorted" RSL scenario of Sec. II. In other words, the curved extra-dimension ψ should appear in the theory in the form of a scalar Brans-Dicke field. Another important ingredient coming from the "distorted" RSL scenario developed here is the fact that the matter degrees of freedom are coupled to the metric induced on the "visible" brane where they live, i.e., $e^{-\psi}g_{ab}$. Finally, it is the hope that such an effective four-dimensional theory would resemble (low-energy) string theory that seems to be the final theory unifying the fundamental interactions.

The effective action for the theory compatible with all of the above requirements is the following

$$S = \int d^4x \sqrt{-g} e^{-\psi} (R - \omega (\nabla\psi)^2 + 16\pi e^{-\psi} \mathcal{L}_m(\lambda, \nabla\lambda, e^{-\psi}g_{ab})), \quad (4.1)$$

where the dilaton field ψ should be identified with the curved extra-dimension, R is the curvature scalar of the metric g_{ab} , the coupling parameter $\omega = \frac{1}{4k^2} - \frac{3}{2}$ and $\mathcal{L}_m$ is the Lagrangian of the ordinary matter fields. It depends on the matter degrees of freedom λ (and their derivatives $\nabla\lambda$) that are directly coupled to the metric $e^{-\psi}g_{ab}$. The dilaton field

ψ is non-minimally coupled both to curvature and to the matter degrees of freedom. Due to the non-minimal coupling of the scalar field to curvature the Newton's constant is not a "true" constant. It varies from point to point like $e^{-\psi}$. In effect, the field equations that are derivable from the action (4.1) are

$$R_{ab} - \frac{1}{2}g_{ab}R = 8\pi e^{-\psi}T_{ab} + \omega[\psi_{,a}\psi_{,b} - \frac{1}{2}g_{ab}(\nabla\psi)^2] + \\ + [-\psi_{||ab} + \psi_{,a}\psi_{,b} + g_{ab}\square\psi - g_{ab}(\nabla\psi)^2], \quad (4.2)$$

where $T_{ab} = \frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g_{ab}}$, $\square\psi = g^{mn}\psi_{||mn}$, and the bar denotes covariant differentiation in a metric sense, i.e., defined through the Christoffel symbols of the metric g_{ab} . Recall that the theory is built over Weyl-integrable geometry so the affine connection does not coincide with the Christoffel symbols. From this equation one sees that Newton's constant $G = e^{-\psi}$. For the dilaton field (literally the curved extra-dimension) the variational principle yields

$$R + \omega(\nabla\psi)^2 - 2\omega\square\psi = -8\pi e^{-\psi}T, \quad (4.3)$$

but, if we take the trace of Eq. (4.2) and combine it with Eq. (4.3) we obtain the following wave equation for the dilaton

$$(2\omega + 3)[\square\psi - (\nabla\psi)^2] = 0. \quad (4.4)$$

The non-minimal coupling of the scalar field to the matter fields allows, in particular, for a variable rest mass m of matter particles in such a way as to preserve the "true" constant character of the dimensionless gravitational coupling constant Gm^2 . In fact, the equation of motion for point-like (uncharged and spinless) particles that is derivable from (4.1) is the following

$$\frac{d^2x^a}{ds^2} + \frac{1}{2}\psi_{,n}\frac{dx^n}{ds}\frac{dx^a}{ds} + \Gamma_{mn}^a\frac{dx^m}{ds}\frac{dx^n}{ds} = 0, \quad (4.5)$$

where, in order to simplify writing, we have dropped the index 4 in the line-element $ds_4 \rightarrow ds$. But we should keep in mind that it is an effective four-dimensional line element that is related to the five-dimensional one through Eq. (2.3). In Eq. (4.5) Γ_{bc}^a is the affine connection of the

four-dimensional spacetime of Weyl-integrable effective configuration. In Sec. III we showed that this equation can be put in the form of a geodesic over Weyl-integrable spacetime (compare with Eq. (3.5) and (3.8))

$$\frac{dP^a}{ds} + m\Gamma_{mn}^a \frac{dx^m}{ds} \frac{dx^n}{ds} = 0, \quad (4.6)$$

where $P^a = m \frac{dx^a}{ds}$ and the mass of the point-like particle depends on the spacetime point through the extra-coordinate ψ ; $m = e^{\frac{\psi}{2}} m_0$. Consequently, $Gm^2 = m_0^2 = \text{const.}$ It is an edifying result: our effective theory respects the strong equivalence principle (SEP). In fact, although the Newton's constant G varies over the four-dimensional spacetime manifold, the units of mass vary in such a way that no local changes in the laws of physics (including the laws of gravity) are "felt". The weak equivalence principle (WEP) is also preserved since particle's mass variations are the same for all particles living on the same brane.

The dynamic equations for the matter degrees of freedom that are derivable from the action (4.1) are of the form

$$T_{||n}^{an} + \frac{1}{2}\psi^{,a}T = 0. \quad (4.7)$$

These can be written in terms of the affine covariant derivative of the Weyl-integrable structure as

$$T_{;n}^{an} + 3\psi_{,n}T^{an} = 0. \quad (4.8)$$

The second term in the left-hand side of Eq. (4.8), i.e., the term that "breaks" vanishing of the affine covariant derivative, originates from variation of the units of measure that is inherent to spacetimes of Weyl-integrable configuration. It may be avoided by allowing the units of measure to vary in such a way that $T^{ab} \rightarrow e^{-3\psi}T^{ab}$. Otherwise it may be viewed as a signal from the extra-dimension. In fact, the dynamic equation of matter in the bulk spacetime is $T_{;N}^{AN} = 0$. It leads, in particular, that $T_{;N}^{aN} = 0$ implies $T_{;n}^{an} + T_{;\psi}^{a4} = 0$. In other words, this equation takes into account possible interchanges of energy between the bulk and the branes through particles on the branes that can escape to the bulk or particles

propagating in the bulk that can enter the branes being trapped there. This situation is analogous to that taking place with point-like particles and that we have explained above in this section. In this last case variations of mass may be viewed as a signal from the extra-dimension (for a detailed analysis of a similar situation see reference [14]).

The strongest constraint on any given theory of gravity is that it must contain general relativity as some limit or particular case. The present theory is a "generalization" of general relativity in the sense that it is a higher (Weyl)-symmetric description of gravity. At the same time Weyl-integrable geometry is a generalization of Riemann geometry which shows higher (conformal) degree of symmetry. Consequently, general relativity together with its geometrical scenario (Riemann geometry) may be recovered from the present description by breaking of the Weyl-symmetry. Symmetry breaking may be implemented either continuously or spontaneously. In the first case it is expected that, some solutions of the wave equation (4.4) allow the dilaton field to evolve from an initial asymptotic state into another asymptotic state, meaning that Weyl symmetry is continuously and asymptotically broken. The second case may be realized by introducing a symmetry breaking self-interaction potential for the dilaton in Eq. (4.1), for instance, of the form $S_{SB} = \lambda \int d^4x \sqrt{-g} e^{-\psi} (\psi^2 - \psi_0^2)^2$. In this case the original Weyl-symmetry of the theory is spontaneously broken by the presence of a "true" constant dimensional value ψ_0^2 .

The present formulation, besides, represents a generalization of general relativity in the sense that it is embeddable within the higher (five)-dimensional (distorted) Randall-Sundrum scenario. In this sense breaking of the original Weyl-symmetry causes brane geometry to be stabilized. In fact, once a given constant value of ψ is reached (either spontaneously or continuously) the location of the brane along the extra-coordinate is fixed in respect to the reference (Planck) brane located at $\psi = 0$. In other words, brane stabilization is a direct consequence of conformal symmetry breaking. Consequently, studying four-dimensional effective gravity theories (built following a "four-dimensional way of thinking") that are embeddable in higher-dimensional scenarios, may be relevant for the study of some issues that are inherent to these scenarios.

V. WEYL-SYMMETRIC EFFECTIVE THEORY AND SPACETIME SINGULARITIES

In this section we shall study one of the more disturbing problems of classical gravity: the singularity problem. We shall see how the scenario developed in the former sections contributes to a possible resolution of this problem.

The "distorted" RSL scenario proposed here is intended to explain physics somewhere after Planck epoch and before conformal symmetry is broken. This way, the early universe may be accommodate well in this model. Recall statement about strong evidence that the early universe underwent a phase where it was five-dimensional [4,5].

However, we shall go a step further and we shall conjecture that, even in the present epoch of the evolution of the universe when branes should be stabilized and conformal symmetry should be broken on cosmological scales, there are sites in our universe (presumably near of gigantic concentrations of energy) where conformal symmetry is yet an exact symmetry of the laws of physics. In this sense, we hope, that our distorted RSL scenario may change the standard black hole picture. With this in mind we shall first study the local problem for the static, spherically symmetric case and then the cosmological situation for Friedmann-Robertson-Walker (FRW) universe filled with a barotropic perfect fluid.

A. The Schwarzschild solution

There is an interesting suggestion in the literature that the Schwarzschild black hole may be considered as a "window" to the extra-dimensions by accommodating the static, spherically symmetric situation within the frame of Kaluza-Klein theory [15]. Inspired by that work we shall consider a similar situation in the frame of our five-dimensional scenario by studying the corresponding solutions to the four-dimensional effective theory. Consequently, we shall consider that there are not any matter degrees of freedom on the "visible brane", i.e., $T_{ab} = 0$ in Eq. (4.2) (literally the vacuum case). We shall use Schwarzschild coordinates

t, r, θ, ϕ , and, considering spherical symmetry we shall search for static solutions to equations (4.2) and (4.4). The corresponding solutions are of Agnese-LaCamera-type [16,17]. In particular, a solution to Eq. (4.4) is found to be

$$\psi - \psi_0 = -q \ln(1 - \frac{2m}{pr}), \quad (5.1)$$

where ψ_0 is an arbitrary integration constant, p and q are arbitrary parameters fulfilling the constraint equation

$$p^2 + (2\omega + 3)q^2 = 1, \quad (5.2)$$

and $\omega = \frac{1}{4k^2} - \frac{3}{2} \geq -\frac{3}{2}$. The solution to Eq. (4.2) yields the following four-dimensional line-element [17]

$$ds_4^2 = -(1 - \frac{2m}{pr})^{p-q} dt^2 + (1 - \frac{2m}{pr})^{-p-q} dr^2 + \rho^2 d\Omega^2, \quad (5.3)$$

where $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$ and the proper radial coordinate $\rho = r(1 - \frac{2m}{pr})^{\frac{1-p-q}{2}}$ has been defined. The curvature scalar of the four-dimensional metric is given as follows

$$R = \frac{4\omega m^2 q^2}{p^2 r^4} (1 - \frac{2m}{pr})^{p+q-2}. \quad (5.4)$$

For a detailed analysis of this solution see Ref. [17]. The definition for the proper radial coordinate leads to the Schwarzschild radial coordinate being constrained: $r \geq \frac{2m}{p}$. From Eq. (5.4) we see that, for $p+q \geq 2$, there is not any spacetime singularity within the allowed range of the Schwarzschild radial coordinate. This solution represents two asymptotic spatial infinities at $r \rightarrow \frac{2m}{p}$ and $r \rightarrow \infty$ respectively that are joined by a wormhole of throat radius $r = (1 + p + q)\frac{m}{p}$ or the invariant surface given by $\rho_{min} = (\frac{p+q+1}{p+q-1})^{\frac{p+q}{2}} \sqrt{(p+q)^2 - 1} \frac{m}{p}$. From Eq. (5.1) it can be seen that $r \rightarrow \frac{2m}{p}$ implies $\psi \rightarrow \infty$, while $r \rightarrow \infty$ yields $\psi \rightarrow \psi_0$. Geodesic completeness of the above solution has been studied in Ref. [17].

Our four-dimensional wormhole solution may be viewed in the higher-dimensional "distorted" RSL scenario as a "five-dimensional" wormhole joining two asymptotically flat branes, one with extra-coordinate $\psi = \psi_0$ and the other at an infinite distance away from

this one. Falling into one such wormhole configuration one can travel in the fifth dimension without "feeling" it. In fact, the geodesic of a point-like particle trapped in the "visible" brane lies all the time on a four-dimensional hypersurface properly bent as to provide such a static five-dimensional configuration.

In this case Weyl symmetry is asymptotically broken on each of the asymptotic branes at $\psi = \psi_0$ and $\psi = \infty$.

B. Friedmann-Robertson-Walker cosmology

Now we shall apply the distorted RSL scenario and the Weyl-invariant effective theory developed here to the particular case of a four-dimensional flat FRW universe filled with a barotropic perfect fluid obeying the equation of state $p = (\gamma - 1)\mu$, where μ is the energy density of the fluid and the barotropic index $0 \leq \gamma \leq 2$. In this case equations (4.2,4.4) and the dynamic equation (4.8) may be simplified by using the conformal technique. We first perform the conformal transformations of the metric

$$\bar{g}_{ab} = e^{-\psi} g_{ab}, \quad (5.5)$$

yielding that the wave equation (4.4) can be simplified to $\ddot{\psi} + 3\bar{H}\dot{\psi} = 0$, ($\bar{H} = \frac{\dot{\bar{a}}}{\bar{a}}$), where $\bar{a}$ is the conformal scale factor $\bar{a} = e^{-\frac{\psi}{2}}a$ and the overdot accounts for derivative in respect to the conformal time $\bar{t} = \int dt e^{-\frac{\psi}{2}}$. Integrating once this equation one obtains

$$\dot{\psi} = \pm \frac{C_1}{\bar{a}^3}, \quad (5.6)$$

where C_1 is an arbitrary integration constant. Integration of the dynamic equation conformal to Eq. (4.8) yields that the conformal energy density $\bar{\mu} = e^{2\psi}\mu = \frac{3}{8\pi} \frac{M}{\bar{a}^{3\gamma}}$, where M is another integration constant. The equation for determining the conformal scale factor can be written as

$$(\bar{a}^3 \bar{H})^2 = M \bar{a}^{3(2-\gamma)} + \frac{C_1^2}{24k^2}. \quad (5.7)$$

If one introduces a new time coordinate $d\bar{t} = \frac{\bar{a}^{3\gamma-3}}{6-3\gamma}d\eta$, then Eq. (5.7) can be readily integrated to yield

$$\bar{a}(\eta) = [\frac{M}{4}(\eta^2 - \eta_0^2)]^{\frac{\alpha}{3}}, \quad (5.8)$$

where $\alpha \equiv \frac{1}{2-\gamma}$ and $\eta_0 = \frac{C_1}{\sqrt{6}kM}$. By integrating Eq. (5.6) one finds

$$\psi^\pm(\eta) = \psi_0 \pm \sqrt{\frac{8}{3}}k\alpha \ln\left(\frac{\eta - \eta_0}{\eta + \eta_0}\right). \quad (5.9)$$

Now we should transform back our solution to the original frame to obtain for the scale factor the following behavior

$$a^\pm(\eta) = \left(\frac{M}{4}\right)^{\frac{\alpha}{3}} e^{\frac{\psi_0}{2}} [(\eta - \eta_0)^{1 \pm \sqrt{6}k} (\eta + \eta_0)^{1 \mp \sqrt{6}k}]^{\frac{\alpha}{3}}. \quad (5.10)$$

This solution shows two branches that are given by the choice of the "+" or "-" signs in Eq. (5.10). The four-dimensional Ricci scalar can be written as

$$R^\pm = -\left[\left(\frac{4}{Me^{\frac{\psi_0}{2\alpha}}}\right)^2 \frac{(\eta - \eta_0)^{\mp \sqrt{\frac{8}{3}}k-2}}{(\eta + \eta_0)^{\mp \sqrt{\frac{8}{3}}k+2}}\right]^\alpha \left[C_1^2 + \frac{3(3\gamma - 4)M^2}{4}(\eta^2 - \eta_0^2)\right]. \quad (5.11)$$

From this we see that the "+" branch is always singular at $\eta = \eta_0$, i.e., for any k there is a cosmological singularity at the beginning of the conformal time coordinate. On the contrary, the "-" branch is singularity-free if $k \geq \sqrt{\frac{3}{2}}$ ($-\frac{3}{2} \geq \omega \geq -\frac{4}{3}$). In this case the geometrical structure of the four-dimensional spacetime is a cosmological wormhole (see Ref. [18] for details). The cosmological time for this branch of the solution is given through the conformal time η as

$$t^- = \frac{\alpha e^{\frac{\psi_0}{2}}}{3} \left(\frac{M}{4}\right)^{\alpha(\gamma-1)} \int d\eta \left[\frac{(\eta - \eta_0)^{\gamma-1-\sqrt{\frac{2}{3}}k}}{(\eta + \eta_0)^{1-\gamma-\sqrt{\frac{2}{3}}k}}\right]^\alpha, \quad (5.12)$$

so, when $\eta \rightarrow \eta_0$, $t^- \rightarrow -\infty$ while $\eta \rightarrow +\infty$ yields $t^- \rightarrow +\infty$, i.e., the cosmological time $-\infty \leq t^- \leq +\infty$. This means that the dilaton (the curved extra-coordinate in our distorted RSL scenario) evolves, in this case, from $\psi^- \sim \infty$ at $t^- \sim -\infty$ into $\psi^- \sim \psi_0$ at $t^- \sim +\infty$. In our brane scenario this means that, in the distant past, we had an infinite separation of the "visible" and "Planck" branes in the extra-dimension (infinite volume

of the extra-dimension). The position ψ of the "visible" brane (recall that the Planck brane is the reference brane) evolves asymptotically into a final finite constant value ψ_0 at the infinite future. In other words, the brane is stabilized at ψ_0 asymptotically. In this case conformal (Weyl) symmetry is continuously broken and standard general relativity over Riemann geometry is recovered asymptotically.

Since our model pretends explaining the evolution of the four-dimensional effective world which we live in before Weyl symmetry breaks down, the "infinite future" $t^- \rightarrow +\infty$ should be understood as that time when conformal symmetry breaking took place.

VI. CONCLUSIONS

In the present paper we developed a four-dimensional effective description of the laws of gravity that embeds naturally in a brane set-up that is a "distortion" of the known Randall-Sundrum scenario. This model contains ingredients that are inherent to the unification scheme of the fundamental interactions. First, it is embeddable in a higher (five)-dimensional spacetime structure. In particular, the extra-coordinate appears in the effective theory in the form of a dilaton field. Geometrically it shows up in a distortion of the usual Riemann geometry into Weyl-integrable geometry over the four-dimensional spacetime manifold. Therefore the extra-coordinate affects the other laws of physics through variations of dimensional scales, for instance, variable masses of point-like particles, variable energy density, etc. Second, the present scenario incorporates the symmetry breaking mechanism. Our four-dimensional effective theory is a Weyl invariant description of the laws of gravity. Standard general relativity can be recovered from it by breaking of this symmetry.

In general, our formulation, is intended to describe the physics over a restricted time interval (possibly after the Planck era and before the conformal symmetry breaking era). However it is argued that, even at present, when conformal symmetry is broken on cosmological scales, there are sites in our universe where conformal symmetry may be an exact symmetry of the laws of physics. In particular, near of gigantic concentrations of energy

(near of the gigantic black hole in the center of our galaxy), the "visible" brane which we live in, may appear distorted or bent by sensible curvature effects in the five-dimensional bulk and conformal symmetry may be present. Falling into one of these sites of gigantic energy concentration means, effectively, traveling in the extra-dimension. However, we never "feel" the extra-dimension since, our geodesic lies always on a four-dimensional hypersurface properly bent as to provide the structure of a static five-dimensional wormhole joining two asymptotically flat branes. In other words, we have just a brane with a non-trivial topological structure embedded in the five-dimensional bulk.

Our four-dimensional description may be relevant for the knowledge of the higher-dimensional structure. In fact, since the extra-dimension appears in our formulation in the form of a dilaton field fulfilling a given wave equation, the corresponding (four-dimensional) solutions contain information about the extra-dimension. In particular, the problem of brane stabilization in essentially five-dimensional formulations (as the original Randall-Sundrum model), being naturally linked with Weyl symmetry breaking, may be explicitly treated within the context of our effective (four-dimensional) description. In this sense, it is of interest, to extend the present formulation to include spontaneous breaking of the Weyl symmetry through inclusion of an appropriate symmetry breaking self-interaction potential for the dilaton (extra-coordinate) in the effective action of the theory.

One of the most interesting features of the present set-up is that it respects both the weak equivalence principle and the strong equivalence principle.

This approach may appear to be less general than postulating from the beginning a five-dimensional theory of gravity. However we think there are, at least, two reasons why it is edifying to elaborate on it. First, it represents a viable alternative to search for a four-dimensional theory consistent with higher dimensions based on a four-dimensional way of thinking. Second, searching for an effective four-dimensional theory pretending to represent a correct description of a higher-dimensional physics means avoiding additional complications related with boundary conditions in five-dimensional brane world scenarios. In particular the field equations may appear to be more familiar and more tractable.

VII. ACKNOWLEDGMENTS

We acknowledge the MES of Cuba for support of this research.

REFERENCES

- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B**429**, 263(1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B**436**, 257(1998).
- [2] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690(1999).
- [3] P. Horava and E. Witten, Nucl. Phys. B**475**, 94 (1996).
- [4] E. Witten, Nucl. Phys. B **471**, 135 (1996).
- [5] J. E. Lidsey, D. Wands and E. J. Copeland, Phys. Rept. **337**, 343(2000).
- [6] F. Darabi and P. S. Wesson, gr-qc/0003045.
- [7] W. Drechsler and H. Tann, Found. Phys. **29**, 1023(1999); W. Drechsler, Found. Phys. **29**, 1327(1999).
- [8] C. Brans and R. H. Dicke, Phys. Rev. **124**(1961)925.
- [9] M. Novello, L. A. R. Oliveira and J. M. Salim, Int. J. Mod. Phys. D, Vol. 1, Nos 3 & 4 (1993)641.
- [10] P. Binétruy, C. Deffayet and D. Langlois, Nucl. Phys. B**565**, 269 (2000); P. Binétruy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. B**477**, 285 (2000).
- [11] J. Lykken and L. Randall, JHEP **0006**, 014(2000).
- [12] P. Binétruy, C. Deffayet and D. Langlois, hep-th/0101234.
- [13] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B**125**, 139 (1983).
- [14] H. Liu and B. Mashhoon, Phys. Lett. A**272**, 26(2000).
- [15] A. Tomimatsu in H. Sato and T. Nakamura (editors), *Gravitational Collapse and Relativity*, 417(World Scientific, 1986).
- [16] A. Agnese and M. LaCamera, Phys. Rev. D**31**, 1280 (1985).

- [17] I. Quiros, Phys. Rev. D**61**(2000)124026.
- [18] I. Quiros, R. Bonal and R. Cardenas, Phys. Rev. D**62**(2000)044042.